

A Massive Dark Object in the S0 galaxy NGC 4350

E. Pignatelli

SISSA/ISAS, via Beirut 4, I-34014 Trieste, Italy

Abstract. We present a new detection of a Massive Dark Object in the S0 Galaxy NGC 4350, obtained applying a new dynamical model on ground-based photometric and kinematic data already present in literature.

There is increasing evidence that most galaxies with a large spheroidal component host a Massive Dark Object (MDO) in their centre, with masses ranging from $\sim 10^8 M_\odot$ to $2 \cdot 10^{10} M_\odot$ (Ho, 1999).

The goal of this work is to investigate the possible presence of an MDO in the nucleus of the S0 galaxy NGC 4350, by applying a new dynamical model (Pignatelli & Galletta, 1999) to ground-based data already present in literature (Seifert and Scorza, 1996; Fisher, 1997). The galaxy was selected on the basis of the high observed rotational velocity of the nuclear disk.

We show that the photometric profiles and the kinematics along the major and minor axes, including the h_3 and h_4 profiles, imply the presence of a central MDO of mass $M_{\text{MDO}} \sim 1.5 - 9.7 \cdot 10^8 M_\odot$, i.e. 0.3–2.8% of the mass derived for the stellar spheroidal component. Models without MDO are unable to reproduce the kinematic properties of the inner stars and of the rapidly rotating nuclear gas.

It is interesting to compare the value we found with the known relations between the masses of the central black holes and the observed “central” stel-




Figure 1. Stellar (*filled symbols*) and gas (*open symbols*) rotational velocities (Fisher, 1997) for the major axis, compared with our best-fit model circular and velocities with (*solid line*) and without (*long-dashed line*) a central MDO. The squares and triangles represent data respectively from the approaching and the receding side.




Figure 2. Comparison between the values of σ_e and M_{MDO} we found for NGC 4350 (filled circle) and the relation given by Gebhardt et al. (2000; left plot) and Ferrarese & Merritt (2000; right plot).

lar velocity dispersions (see Fig. 2). While the best-fit given by the stellar kinematics is almost 10 times higher than the average relation, the lower limit we find is consistent (within the observed scatter) with this relation. On the other hand, we expect (because of selection effects) this galaxy to host a MDO a bit more massive than the average for this velocity dispersion. Moreover, it is known that ground-based stellar kinematics -if the systematic errors are not properly taken into account- gives M_{MDO} higher than the ones obtained with HST spectroscopy. In this work, when all the possible sources of errors are taken into account, we find results which are again consistent with the HST average values (see Pignatelli et al., 2000 for a complete discussion).

We also found that the stellar population of this galaxy comprise of an exponential disc (27% of the light) and a diffuse spheroidal component (73% of the light) that cannot be represented by a simple de Vaucouleurs profile at any radius. The M/L ratios we found for the stellar components (respectively 3.3 and 6.6) are typical of those of disc and elliptical galaxies.

References

- Ferrarese, L. & Merritt, D. 2000, *ApJ*, 539, L9
- Fisher, D. 1997, *AJ*, 113, 950
- Gebhardt, K., Bender, R., Bower, G., Dressler, A. et al. 2000, *ApJ* 539, L13
- Ho, L. 1999, in *Observational Evidence for the Black Holes in the Universe*, Conference held in Calcutta, January 11-17th, 1998, 157
- Pignatelli, E. & Galletta, G. 1999, *A&A*, 349, 369
- Pignatelli, E., Salucci, P., Danese, L. 2000, *MNRAS*, in press (astro-ph/0008429)